

GBlock® gene fragments

G-block gene fragments can be directly ordered from idtdna.com

List of Gblocks®

Virus	Gblock® sequences
HAdV	ATGATGCCGCAATGGTCTTACATGCACATCGCCGGCAGGACGCCTCGGAGTATCTGAGCCCGGGCACACACA CACACCTGGTGCAATTTGCCCGGCCACCGATACGTACTTCAGCCTGGGGAACAAGTTCAGAAATCCCGCTGC GATTCGTGCCAGTCGACCGCGAGGACACCGCTTATTCTTACAAAGTGCCTTTACGCTGGCCGTGGGCGACAA CCGGGTGTTGGACATGGCCAGCACCTACTTTGACATCCGCGGCGTGCTGGATCG
JCPyV VP1 region and IG region	TTCTCCCTATTAGCACTTTGTCATTTTACTGCTTTTGCAGCAAAAAATTACTGCAAAAAAGGGAAAAACAAG GGAATTTCCCTGGCCTCTAAAAAGCCTCCACGCCCTTACTACTTCTGAGTAAGCTTGGAGGCGGAGGCGGCC TCGGCCTCTCACACAAAAAAGGGAAGGGATGAGCCAGAGCTGTTTTGGCTTGTACCAGCTGGCCAT GGTCTTCGCCAGCTGTCAGTAAGGCTTCTGTGAAAGTTAGTAAACCTGGAGTGGAATAAAAAAGAGCT CAAAAAGACAGAGACACAGTGGTTTGAATTGCGAAAGTCTTGGGTCTTCTACCTTTTTCACACACATTTCT TTTTAGGTGGGGTAGAGTGTGGGATCCTGTGTTTTCATCATCTGGCAAACATTTGAATTCGTTGAATGTTG GGTTCCTGATCCCACCAGAAATGAAAACACAAGATATTTGGGACACTAACAGGAGGAGAAAATGTTCTCCA GTTCTTCATATAACAAACTGCCACAACAGTGTGGCTTGATGAATTTGGTGTGAAAGCCACACATGTGAA AAGTGCAGTTTTCTGTGTCTGCACCAGAGGCTTCTGAGACCTGGGAAAAGCATTGTGATTGTGATTGAGT GCTTGATCCATGTCCAGAGTCTTCTGCTTCAAGATCTTCTCTAGGAAAGTCAAGAATGGGTCTCCC
PAdV and OPyV	TCCCCGGATGTCATGGACAACGTCAACCCTTCAACCACCACCGCTCCTGGGCAACGGCCGCTACTGCAAGTT CCACATCCAGGTGCCGCAAAAGTTCTTTGCCCTCAAGAGCCTGCTGCTCCTGCCGGGGCGACCGGCCAACGT CACCAACTGCCCATCTCCATTCCAGCCGCAACTGGCCGCATCAGCATCCAGTTCGACTCCTCCGTGGCCTGC ACACCCATGGACACAGAGATGGCCTCTTTCTCACAGCTGCAGACATTGTGGGCATGATTACCAAGCCAGACAG TGGGCACATGGCTTTTCGTGGGCTCCCAAGGTAATCAATGTCACTTAAGAAAGAGAGTGGTTAAGAACCA TATCCTGTTAGCACATTGCTGAATCTCTTTTCAAATCTTATGCCAGATTGGAAGGCCAGGCCATGACTGG AGATAAATCC
BPyV and Ch/TyPV	GAAAAGGTATTCGCCCTCTGCTGGTCAAGGCTCAGCAATATGTGAGGATTTCAAAGCCCCCTATCATCCCACAG TACCCAGGTGCTAGATCCTACCCTCAAGGGAATCTAGACAAAGATGGTGTGTATCCTGTTGAGTGTGGTGTC CAGATCCAAGTAAAAATGAACCGCTTCTGATGAGAATGGCGTAGGCCCGCAGACTGGTGAAGAACCATACC CCATTGCCAGCATACTCACACCCATGGACACGTAATTGGAATTGTGATACTATATGGGCTGAAAATAAGCCG CCATTGTGTCTGCTTATGCGTGACATGGATCAGCGTGCACTTTAGTCCACGAGATTGGCAACAAGTAAAT AATTATTCGAGATGGCGCCACGCGCCACACACAGTACACGTGAAAATCTTTAACCTGCAAGTCATCCAGAA AACCGTAACAGATAGCTGACAGGAACAATCCAAATATTTGCAGATCAGGAAGG
MS2	CATAAAGGTGCCTACAAGCGAAGTGGGTGCATCGTGGGGTCCCGTACGAGGAGAAAAGCCGGTTTCGGCTTC TCCCTCGACGCACGCTCCTGCTACAGCCTTCCCTGTAAGCCAAAATTGACTTACATCGAAGTGCCGCGAGAA CGTTGCGAACCGGGCGTCGACCGAAGTCTGCAAAAGGTCACCCAGGGTAATTTAACCTTGGTGTGCTTTA GCAGAGGCCAGGTCGACAGCCTCACAACCTCGCGACGCAAAACCATTGCGCTCGTGAAGGCGTACTGCGCCTC GTCGCGTAATTGGCGCCAGGCGCTCCGCTACCTTGCCCTAAACGAAGATTA

Gblocks® were designed with small deletions to detect cross-contamination. G-block amplicons have a different length than nPCR amplicons (see amplicon length of the virus or synthetic GBlock® PCR and nPCR fragments).

List of primers and probes for each specific PCR, nPCR and qPCR.

PCR and nPCR primers (5' - 3')	qPCR primers and probes (5' - 3')
HAdV^{1,2}	
Hex1deg: GCCSCARTGGKWCWTACATGCACAT	AdF:CWTACATGCACATCKCSGG
Hex2deg:CAGCACSCCYCGRATGTCAAA	AdR:CRGGGCRAAYTGCACCAG
Nehex3deg:GCCCGYGCMACYGAYACSTACTTC	AdP:6-FAM-CCGGGCTCAGGTACTCCGAGGCGTCT-BHQ1
Nehex4deg: CCYACRGCCAGYGTRWAYCGMRCYTTGTA	
JCPyV^{3,4,5}	
VP1 region	JE3F: GGAAAGTCTTTAGGGTCTTCTACCTT
JR1: CCCTATTCAGCACTTTGTCC	JE3R: ATGTTTGCCAGTGATGATGAAAA
JR2: CAAACCACTGTGTCTCTGTCT	JE3P: 6-FAM-AGGATCCCAACTCTACCCACCTAAAAAGA-BHQ1
JR3: GGGAAATTTCCCTGGCCTCCT	
JR4: ACTTTACAGAAGCCTTACG	
IG region	
EP1A: TGAATGTTGGGTTCTGATCCCACC	
EP2A: ACCATTCTTGACTTTCTAGAGAG	
P1A: CAAGATATTTGGGACACTAACAGG	
P2A: CCATGTCCAGAGTCTTCTGCTTCAG	
PAdV^{6,7}	
PALF: GATGTCATGGACAACGTCAAC	Q-PAdV-F AACGGCCGCTACTGCAAG
PARF: CACGGAGGAGTCGAACTGGATG	Q-PAdV-R AGCAGCAGGCTCTTGAGG
PALN: TACTGCAAGTTCCACATCCAGGT	Q-PAdV-P 6-FAM-CACATCCAGGTGCCGC-BHQ1
PARN: GGAATGGAGATGGGCAGGTT	
OPyV⁸	
Ov_FI: AGATGGCCTCTTTCTACA	Ov_qFI: TTCTCACAGCTGCAGACATTG
Ov_RI: TTTATCTCCAGTCATGGCC	Ov_qRI: TCCAATCTGGGCATAAGATTTG
Ov_nFI: AGACATTGTGGGCATGATTA	Ov_qPI: 6-FAM-ATGATTACCAAGCCAGACAGTGGG-BHQ1
Ov_nRI: TTCCAATCTGGGCATAAGAT	
BPyV^{6,9}	
VP1F: GGTATTCGCCCTCTGCTGGTCAAG	QB-F1-1: CTAGATCCTACCCTCAAGGGAAT
VP1R: GCTGGCAATGGGGTATGGGTTCT	QB-R1-1: TTA CTGGATCTGGACACCAAC
VP2F: ATTTCAAAGCCCCCTATCATC	QB-P1-2: 6-FAM-GACAAAGATGGTGTGTATCCTGTTGA-BHQ1
VP2R: GCCTACGCCATTCTCATCAAG	
Ch/TyPV¹⁰	
VP2-Par1: TGGAATTGTGATACTATATGGG	Q-PaV-F AGTCCACGAGATTGGCAACA
VP2-Par2: TCYTGATCTGCAAATATTTG	Q-PaV-R GCAGGTTAAAGATTTTACG
VP2-Par3: CATTGTGTCTGTCTWATGCGTGAC	Q-PaV-Pr 6-FAM-AATTATTCGAGATGGCGCCACG-BHQ1
VP2-Par4: GTTTTCTGGATGACTTGCA	
MS2¹¹	
-	pecson-2F: AAGGTGCCTACAAGCGAAGT
	pecson-2R: TTCGTTTAGGGCAAGGTAGC
	PecP-2: 6-FAM-ATCGTGGGGTCGCCGTACG-BHQ1

Amplicon length of the virus or synthetic GBlock® PCR and nPCR fragments

Virus	Amplicon length (bp)		Gblock® amplicón length (bp)	
	PCR	nPCR	PCR	nPCR
HAdV	313	173	263	123
JCPyV				
VP1 region	736	667	314	186
IG region	587	387	293	223
PAdV	612	344	209	108
BPyV	527	263	227	148
OPyV	227	162	211	163
Ch/TyPV	372	251	231	149

References:

1. Allard A., Albinsson B., Wadell G. Rapid typing of human adenoviruses by a general PCR combined with restriction endonuclease analysis. *J. Clin. Microbiol.* 2001. 39:498-505.
2. Hernroth BE., Conden-Hansson AC., Rehnstam-Holm AS., Girones R., Allard AK. Environmental factors influencing human viral pathogens and their potential indicator organisms in the blue mussel, *Mytilus edulis*: the first Scandinavian report. *Appl. Environ. Microbiol.* 2002. 68: 4523-33.
3. Bofill-Mas S., Albinana-Gimenez N., Clemente-Casares P., Hundesa A., Rodriguez-Manzano J., Allard A., Calvo M., Girones R. Quantification and stability of human adenoviruses and polyomavirus JCPyV in wastewater matrices. *Appl. Environ. Microbiol.* 2006. 72: 7894-7896.
4. Bofill-Mas S., Formiga-Cruz M., Clemente-Casares P., Calafell F., Girones R. Potential transmission of human polyomaviruses through the gastrointestinal tract after exposure to virions or viral DNA. *J Virol* 2001. 75(21):10290-9.
5. Pal A., Sirota L., Maudru T., Peden K., Lewis AM. Real-time PCR assays for the detection of virus-specific DNA in samples with mixed populations of polyomaviruses. *J. Virol. Methods* 2006. 135(1):32-42.
6. Maluquer de Motes C., Clemente-Casares P., Hundesa A., Martin M., and Girones R. 2004. Detection of bovine and porcine adenoviruses for tracing the source of fecal contamination. *Appl. Environ. Microbiol.* 70:1448-1454.

7. Hundesa A., Maluquer de Motes C., Albinana-Gimenez N., Rodriguez-Manzano J., Bofill-Mas S., Suñen E., Girones R. Development of a qPCR assay for the quantification of porcine adenoviruses as an MST tool for swine fecal contamination in the environment. *J Virol Methods* 2009. 158(1-2):130-135.
8. Rusiñol M., Carratalà A., Hundesa A., Bach A., Kern A., Vantarakis A., Girones R., Bofill-Mas S.. Description of a novel viral tool to identify and quantify ovine faecal pollution in the environment. *Sc Total Env* 2013. 458-460C:355-360.
9. Hundesa A., Bofill-Mas S., Maluquer de Motes C., Rodriguez-Manzano J., Bach A., Casas M., Girones R. Development of a quantitative PCR assay for the quantitation of bovine polyomavirus as a microbial source-tracking tool. *J Virol. Methods.* 2010. Feb 163(2):385-389.
10. Carratalà A., Rusiñol M., Hundesa A., Biarnés M., Rodriguez-Manzano J., Vantarakis A., Kern A., Suñen E., Girones R., Bofill-Mas S. A novel tool to trace poultry fecal contamination in the environment: specific detection and quantification of chicken/turkey parvoviruses. *Appl. Environ. Microbiol.* 2012. 78(20):7496-9
11. Pecson BM., Martin LV., Kohn T. Quantitative PCR for determining the infectivity of bacteriophage MS2 upon inactivation by heat, UV-B Radiation, and singlet oxygen: advantages and limitations of an enzymatic treatment to reduce false-positive results.. *Appl. Environ. Microbiol.* Sept. 2009, p. 5544-5554-33.